

# A probabilistic view on possibility distributions\*

## Christer Carlsson

IAMSR, Åbo Akademi University,  
Lemminkäinenengatan 14B, FIN-20520 Åbo, Finland  
e-mail: christer.carlsson@abo.fi

## Robert Fullér

Department of Operations Research, Eötvös Loránd University,  
Pázmány Péter sétány 1C, H-1117 Budapest, Hungary  
e-mail: rfuller@mail.abo.fi

## Péter Majlender

IAMSR, Åbo Akademi University,  
Lemminkäinenengatan 14B, FIN-20520 Åbo, Finland  
e-mail: christer.carlsson@abo.fi

*Abstract:* In this work we shall give a probabilistic interpretation of possibilistic expected value, variance, covariance and correlation.

## 1 Probability and possibility

In probability theory, the dependency between two *random variables* can be characterized through their joint probability density function. Namely, if  $X$  and  $Y$  are two random variables with probability density functions  $f_X(x)$  and  $f_Y(y)$ , respectively, then the density function,  $f_{X,Y}(x, y)$ , of their joint random variable  $(X, Y)$ , should satisfy the following properties

$$\int_{\mathbb{R}} f_{X,Y}(x, t) dt = f_X(x), \quad \int_{\mathbb{R}} f_{X,Y}(t, y) dt = f_Y(y)$$

---

\*in: *Proceedings of the Sixth International Symposium of Hungarian Researchers on Computational Intelligence*, Budapest, [ISBN 963 7154 43 4], November 18-19, 2005 57-61

for all  $x, y \in \mathbb{R}$ . Furthermore,  $f_X(x)$  and  $f_Y(y)$  are called the the marginal probability density functions of random variable  $(X, Y)$ .  $X$  and  $Y$  are said to be independent if the relationship  $f_{X,Y}(x, y) = f_X(x)f_Y(y)$  holds for all  $x, y$ . The expected value of random variable  $X$  is defined as

$$E(X) = \int_{\mathbb{R}} x f_X(x) dx.$$

The covariance between two random variables  $X$  and  $Y$  is defined as

$$\text{Cov}(X, Y) = E((X - E(X))(Y - E(Y))) = E(XY) - E(X)E(Y),$$

and if  $X$  and  $Y$  are independent then  $\text{Cov}(X, Y) = 0$ . The variance of random variable  $X$  is defined by  $\sigma_X^2 = E(X^2) - (E(X))^2$ . The correlation coefficient between  $X$  and  $Y$  is defined by

$$\rho(X, Y) = \frac{\text{Cov}(X, Y)}{\sqrt{\text{Var}(X)\text{Var}(Y)}}$$

and it is clear that  $-1 \leq \rho(X, Y) \leq 1$ .

A *fuzzy number*  $A$  is a fuzzy set of the real line with a normal, (fuzzy) convex and continuous membership function of bounded support. The family of fuzzy numbers will be denoted by  $\mathcal{F}$ . Fuzzy numbers can be considered as possibility distributions [2, 5]. If  $A \in \mathcal{F}$  is a fuzzy number and  $x \in \mathbb{R}$  a real number then  $A(x)$  can be interpreted as the degree of possiblity of the statement "x is A". A fuzzy set  $C$  in  $\mathbb{R}^n$  is said to be a joint possibility distribution of fuzzy numbers  $A_i \in \mathcal{F}, i = 1, \dots, n$ , if it satisfies the relationship

$$\max_{x_j \in \mathbb{R}, j \neq i} C(x_1, \dots, x_n) = A_i(x_i)$$

for all  $x_i \in \mathbb{R}, i = 1, \dots, n$ . Furthermore,  $A_i$  is called the  $i$ -th marginal possibility distribution of  $C$ , and the projection of  $C$  on the  $i$ -th axis is  $A_i$  for  $i = 1, \dots, n$ .

Fuzzy numbers  $A_i \in \mathcal{F}, i = 1, \dots, n$  are said to be non-interactive if their joint possibility distribution  $C$  satisfies the relationship

$$C(x_1, \dots, x_n) = \min\{A_1(x_1), \dots, A_n(x_n)\},$$

or, equivalently,  $[C]^\gamma = [A_1]^\gamma \times \dots \times [A_n]^\gamma$  holds for all  $x_1, \dots, x_n \in \mathbb{R}$  and  $\gamma \in [0, 1]$ . Marginal probability distributions are determined from the joint one by the principle of 'falling integrals' and marginal possibility distributions are determined from the joint possibility distribution by the principle of 'falling shadows'.

If  $A, B \in \mathcal{F}$  are non-interactive then their joint membership function is defined by  $A \times B$ , where  $(A \times B)(x, y) = \min\{A(x), B(y)\}$  for any  $x, y \in \mathbb{R}$ . It is clear that in this case any change in the membership function of  $A$  does not effect the second marginal possibility distribution and vice versa. On the other hand,  $A$  and  $B$  are said to be interactive if they can not take their values independently of each other [2].

Let  $A \in \mathcal{F}$  be fuzzy number with  $[A]^\gamma = [a_1(\gamma), a_2(\gamma)]$ ,  $\gamma \in [0, 1]$ . A function  $f: [0, 1] \rightarrow \mathbb{R}$  is said to be a weighting function if  $f$  is non-negative, monoton increasing and satisfies the following normalization condition

$$\int_0^1 f(\gamma) d\gamma = 1.$$

## 2 A probabilistic interpretation of possibilistic expected value, variance, covariance and correlation

The  $f$ -weighted *possibilistic expected value* of  $A \in \mathcal{F}$ , defined in [3], can be written as

$$E_f(A) = \int_0^1 \frac{a_1(\gamma) + a_2(\gamma)}{2} f(\gamma) d\gamma = \int_0^1 E(U_\gamma) f(\gamma) d\gamma,$$

where  $U_\gamma$  is a uniform probability distribution on  $[A]^\gamma$  for all  $\gamma \in [0, 1]$ .

The  $f$ -weighted *possibilistic variance* of  $A \in \mathcal{F}$ , defined in [3], can be written as

$$\text{Var}_f(A) = \int_0^1 \sigma_{U_\gamma}^2 f(\gamma) d\gamma.$$




Figure 1: The case of  $\rho_f(A, B) = 0$ .

The  $f$ -weighted *measure of possibilistic covariance* between  $A, B \in \mathcal{F}$ , (with respect to their joint distribution  $C$ ), defined by [4], can be written as

$$\text{Cov}_f(A, B) = \int_0^1 \text{Cov}(X_\gamma, Y_\gamma) f(\gamma) d\gamma,$$

where  $X_\gamma$  and  $Y_\gamma$  are random variables whose joint distribution is uniform on  $[C]^\gamma$  for all  $\gamma \in [0, 1]$ .




Figure 2: The case of  $\rho_f(A, B) = 1$ .




Figure 3: The case of  $\rho_f(A, B) = -1$ .

The  $f$ -weighted possibilistic correlation of  $A, B \in \mathcal{F}$ , (with respect to their joint distribution  $C$ ), defined in [1], can be written as

$$\rho_f(A, B) = \frac{\int_0^1 \text{cov}(X_\gamma, Y_\gamma) f(\gamma) d\gamma}{\left( \int_0^1 \sigma_{U_\gamma}^2 f(\gamma) d\gamma \right)^{1/2} \left( \int_0^1 \sigma_{V_\gamma}^2 f(\gamma) d\gamma \right)^{1/2}}.$$

where  $V_\gamma$  is a uniform probability distribution on  $[B]^\gamma$ . Thus, the possibilistic correlation represents an average degree to which  $X_\gamma$  and  $Y_\gamma$  are linearly associated as compared to the dispersions of  $U_\gamma$  and  $V_\gamma$ . If  $[C]^\gamma$  is convex for all  $\gamma \in [0, 1]$  then  $-1 \leq \rho_f(A, B) \leq 1$  for any weighting function  $f$  (see [1] for details).

### 3 Examples

First, let us assume that  $A$  and  $B$  are non-interactive, i.e.  $C = A \times B$ . This situation is depicted in Fig. 1. Then  $[C]^\gamma = [A]^\gamma \times [B]^\gamma$  for any  $\gamma \in [0, 1]$  and we have  $\text{Cov}_f(A, B) = 0$  (see [4]) and  $\rho_f(A, B) = 0$  for any weighting function  $f$ . In the case, depicted in Fig. 2, if  $u \in [A]^\gamma$  for some  $u \in \mathbb{R}$  then there exists a unique  $v \in \mathbb{R}$  that  $B$  can take. Furthermore, if  $u$  is moved to the left (right) then the corresponding value (that  $B$  can take) will also move to the left (right). This property can serve as a justification of the principle of (complete positive) correlation of  $A$  and  $B$ . In the case, depicted in Fig. 3, if  $u \in [A]^\gamma$  for some  $u \in \mathbb{R}$  then there exists a unique  $v \in \mathbb{R}$  that  $B$  can take. Furthermore, if  $u$  is moved to the left (right) then the corresponding value (that  $B$  can take) will move to the right (left). This property can serve as a justification of the principle of (complete negative) correlation of  $A$  and  $B$ .

### References

- [1] C. Carlsson, R. Fullér and P. Majlender, On possibilistic correlation, *Fuzzy Sets and Systems*, 155(2005) 425-445.
- [2] D. Dubois and H. Prade, *Possibility Theory: An Approach to Computerized Processing of Uncertainty*, Plenum Press, New York, 1988.
- [3] R. Fullér and P. Majlender, On weighted possibilistic mean and variance of fuzzy numbers, *Fuzzy Sets and Systems*, 136(2003) 363-374.
- [4] R. Fullér and P. Majlender, On interactive fuzzy numbers, *Fuzzy Sets and Systems*, 143(2004) 355-369
- [5] L. A. Zadeh, Concept of a linguistic variable and its application to approximate reasoning, **I, II, III**, *Information Sciences*, 8(1975) 199-249, 301-357; 9(1975) 43-80.